

Chloroform

Chloroform

A case of Erysipelas of the face
at the ninth day, pustules were
seen on the face. The patient is even
drinking an impure water.
Dr. Chapman

A case of Erysipelas - Chloroform in
Bath, bleeding, Chloroform in the
crevice &c

A case of chronic Erysipelas. An
inflammation of the mucous membrane
of the intestine. A new Erysipelas
taken away & removed of blood which
would be purgative. Chloroform over the Ab-
domen small doses of Calomel, Opium
Eucalyptus - as may be required. Chloroform
except small portions of a mucous
layerous nature. Chloroform a new
method. Chloroform in the
face heated in the face
Chloroform in the face
Chloroform in the face

but where the skin is dry we know to the contrary, was ordered
in Enema a tea cup full of
weak tea with 80 drops of Hand-
man

Maria Pott, being affected in
great terror, for which Calomel
and Opium was given. These
symptoms of Pneumonia the
mucous thrown up is thick
transparent and adhering to
the side of the vessel, as the
breath is bloody in some cases
be heard in the cells of the lungs
by means of the stethoscope
as it is getting better it is
again changed, we ought to
know whether the breathing is
carried on with the abdominal
muscles or the thorax, after
the blisters on the side applied
I use a Decoction of Nitro-agric
Lactar Emulic inside in 8 pills
or Calomel but must not be
carried too far

There are inflammation of the stomach
and of the stomach and the intestines
and both are sometimes complicated
with the fever (when it is accom-
panied with diarrhoea)

In this lowest state, Berley water
if you give any skin pills will be
detrimental - But we blight give
small doses of Rochelle Salts
or castor purgative or sometimes not
purge in fever

Riarrhea small doses Castor Oil
and small doses of Ipecac and Opium
with low diet of mucilaginous drinks

Low fever of a typhoid nature
to be treated by evacuation from
the bowels and mild diaphoretics
pain in the abdomen relieved by
cupping, this fever was formerly
treated by acetate of Potash or
any other practice

Intermittent

Syphilis and Cancer. Tomorrow to be
purged to be cured without bark &c
In this stage he would be thrown
into a typhus fever


now given from Colic
pain in the head exc. here and there
much, in this case an affection of the
Mucous surface because the head is
so much affected

If Acute Rheumatic Inflammation
Cochicum a specific remedy
in this disease and ^{Causes} ~~Causes~~ in Intes-
tinal diseases the medicine may
be good and to be given only
until it causes pain or bloating
in stools may be given

Cynchona may be given, Decoc-
tion of Sarsaparilla has cured the
above disease by continuing a
quant daily & a course of
Mercury often of great harm, in
giving the medicine a susceptibility
to a relapse of Rheumatism

In Syphilitic cases given Sarsapa-
rilla with ~~etc~~ of good Corrosive sub-
the medicine being given in too large
doses is the cause of all future trouble

2.
John W.



Gibson

Ulcerative stage, ^{does not effect cure} if the adhesion
then suppuration and then the Ulcera-
tion by a small red spot on the sur-
face and extends itself, not much
pain, does not extend itself inwards
but outwards, sometimes without
suppuration, sometimes ulceration
and gangrene go at the same
time and are sometimes on the same
footings, if it passes this stage then the
disease looks as if for cure, exposure will
promote granulation, granulation
extends, themselves rapidly, and do
with and if not they are not healthy
Care 2 things careful, the action beca-
ried not too far and the other there
be not too much weakness in mis-
takes nothing more than exclude out-
ward affection deeped regularly. bandages
of cotton if no granulation the day
lint. ~~the~~
Gangrene or mortification happens per-
much debilitated and always takes
on the same appearance as from
solation cause, and may be known
by - a general redness not bright red
approaching sometimes to brown, in thin
livid spots there increase while it
a light blue

the pulse frequent tremulous, weak and
if full it is sure death, thirst, urine
voided insensibly, dry gangrene
eating vitiated age, commences the
toe extends to thighs & one case
where the leg of a man 70 years old became
black, one case commencing by a dark
spot on one of the toes cuticle always
early detached when attended with
fever and debility barks &c to be
employed, blood letting but not
too much and with purgatives
dozers powder, bark the best
if cannot be taken try quinine
salt of bark but best of ~~the~~ all
strong wine and good porter,
keep off smell fomenting pou-
tices - if when we called to a limb
badly mangled so that mortifi-
cation must take place we an-
ticipate immediately, if it shall
already mortify we must till
inflammation may be observed on
the edges and then

Of wounds all of Contused Punct
Punct Case &c Contused are such
as are caused velocity by a cannon
ball strike before the strength is
spent it called Contused, a strong
man receive more wound than
a weak one as the muscle does
not give way, a man struck with
a club receive a contused wound
and remain longer than any other
sometimes a contused wound produces
cavous ~~lesions~~

Punctured wound are cuts not so
revise as ~~tent~~ but that the parts come
in contact with each other as soon
as the knife is drawn away —

Penetrating the extent greater than
Punctured & very dangerous for their
depth — Lacerated wounds

produced mechanically &c become
the and contused wounds worst of
all wounds, Incised wound cuts

&c — Union is the first intention
Contused wounds may sometimes may call
Suture Entrement for keeping the parts

are Suture and bandage, stitching
elastic made of akicuin & resin,
Sutura Corana, if
the wounds are very deep like
ought to have the middle
very crooked

The adhesive to be preferred to all
other when it can be used -
Gun shot Wound it is lacerated
contused if the ball strikes
fair it is generally a lacerated
but if ~~strikes~~ glance off it is
a contused which is very diffi-
cult to cure, bone with suffer
more than muscles and maily
more than Cellular, a ball
passing into a dense medium than
when into a rare one, if the
velocity of the ball there is a
tremor over the man and he
does not know he is wounded
but when it is of moderate force
it give more pain, the more the
vitality of a place is wounded
the less will be the pain, a ball
of slight force will sometimes
pass a creature under the skin
which is mark by a red line
Inflammation in Gun shot
wounds as in contused wounds
they do not inflame so soon
as cutting wounds, the greater
the velocity the greater will
be the shock.

If a ball pass through the abdomen
and intestine the patient passes stool
from the Anus regular for 3 or 6
days more or less.

Wounded vessels are seldom closed
is if by a sharp instrument, untill
the patient is put to rest then some
time they bleed, wound sometime
partially heal and then break open
and blood and matter is discharged
and last for a long time with ama-
cration, A man wounded is taken
with a tremor and fear uncon-
mon and even among smaller
ones. Treatment.

1. When the wound first received

2. The consequence of wounds,
The first thing to fear is hemorrhage
must lay open the wound and take
up the vessel immediately and
not put on a tourniquet which
is very dangerous, when hemorrhage
is stopped we put him to bed &
some dispute the propriety of exam-
ining gun shot wounds, when
an examination is necessary
it must be done in

mediately after the accident but
there only when spilled bone or
or other substances carried into
it with other circumstances
such a ball passing the lung liver
&c we ought to be careful if any
necessary to do so - the finger
to be used in preference instrument
if it can be admitted and by
this advice with more certainty
the long gun shot probe the best
or sounder in some circumstances
the best is to dilate the wound or get
it out with scope &c. When the soldier
is carried for a day or two before he is
carried in the hospital, we generally
have much inflammation with a
hard and inflamed ring around
the wound, the hole where the ball
enters ~~the~~ it must be suppurate
the first to be done is to moderate
these symptoms we may employ poultices
and all cure for inflammation
but does not admit blood letting to
well as other causes of inflammation
especially if he shall have lost
much blood - Next to the remove
the cause of suppuration and this
done without violence with
forceps and take care

necessaries of Practicable & the last
sometimes for years with emaciation
sickness and cough & the to the feet
with nourishing food and take
barks and Tonics, need
Wounds of the head from falls or
of children if the blow violent the
the nose rises very great and is because
some blood vessels, have been broken
if this blood should remain for 2 or 3
can weeks the bone becomes carious
if not very large do not make an inci-
sion in it immediately but afterwards
nearly open it take out clotted
blood tie up blood vessels &
new born infants sometimes wash
with spirits and water - ~~Scars~~
in blow on the head cause affection
of the liver, erysipelis of the head
a blister or mercurial, continue
if general system be very much dis-
turbed particula- and to diet
in infants of the head save it clean and
examine very carefully indeed
not examine the bone unless there
are symptoms of very great danger
nervous symptoms follow sometimes
after wounds of the head by affecting
the nerve, this treated by cutting the
nerve or giving juice if practicable

In injuries of the scalp it is
always necessary wash the part
clean and place them to
get them, likewise in cut ear
nose, lips &c be careful to
place them even &c.

Wounds of lower face of which generally die
by commotion of the brain, but of the
chin only not so, of the lower limbs
rather very great and secondary hemo-
rrage very frequent almost impossi-
ble to take up the artery, when the toe
is cut is slight by the teeth place it
in a fig cut in a slit &c
Neck, if cut ~~near~~ very near the chin
it may recover but if cut low
down almost immediate death
if high up not easy to nourish him
Sometimes not cut ~~so~~ so deep
as the wind pipe there are the
interrupted suture on the skin
the support the patient by some
confidence, the best use a com-
mon leather stock around the
neck and leave a hole to dress the
wound, to nourish the patient
it has sometimes been done by passing
a catheter &c into the ~~stomach~~
&c

Gun shot wound of the neck not so dan-
gerous as cut wounds but to be treated
much in the same way, Gun shot
wound generally are thigh, but
~~Dangerous wounds about the~~
~~Clavicle, the Sub Scapular~~
artery wounded sometimes & very
frequently a bullet lies under the
skin always take it out & in
wounds of the chest the most dan-
gerous we meet with by passing
through the lungs but even this does
not always happen balls sometimes
pass ~~under~~ the skin a great way under
the skin and may be traced after two
or three days by a red line of the
skin & all pieces of cloth ought
to be taken, if the lungs &c are wounded
& the have a discharge of blood
from the mouth, if it be the in-
tercostal artery cut down and tie
up the artery! Emp^{ma} is a formida-
ble disease by which the patient is blown
but is off no danger ~~but~~ and requires
two or 3 incision of the scapula only
one remedy for hemorrhage from the
lungs bleed the patient in the arm again
into a vein, and cut the vein, and

If he get over 2 or 3 days he may be
thought in a fair way, the forma-
tion of matter or sinuses in the chest
or lungs which is sometimes thrown out
by the mouth &c. if this wound be
in the chest and thought to be mor-
tal do not introduce the probe &
but depend altogether in the general
treatment, by introducing probe &
you rupture blood vessels &
lungs not ~~only~~ always wounded when
the air enters in and out of the open-
ing but only in the Chest &c.

That if the lungs are wounded the blood
rushes out of the Mouth and almost
chookes him or dies from

If both lungs wounded death almost
immediately - Treatment ~~short~~
bleed ~~the~~ very great deal and close
the external wounds as soon as
possible, if by the symptoms you
know matter is formed make
an opening lower down on the
diaphragm, & give them in rapid
injection and in suppuration stay
nourish the patient as much as pos-
sible with beef steak and port wine
preference to Madeira &c.

Wounds of the head &c

Wounds of the abdomen, which is some-
times does not enter more than beneath
the skin sometimes it is wounded
by a knife and the intestine and peri-
tonium, if an intestine be forced out of
a hole of the wall cut a little and it
will readily receive, to be treated
or less diet take care of inflama-
tion and treat as wounds of the chest

Wounds of the stomach very danger-
ous, never stick the ~~the~~ edges of the sto-
mach, but attach it to the adjoining
part, never search too much for
the wall which sometimes do not as
much harm as the search

Wounds of Peritonium, intestines
Liver &c are almost immediate death
but not quite so and attend to the gene-
ral symptoms &c

Kidneys and Bladder (Almost certain)
death and difficult to cure, also of the
Genital Organs &c

On Wounds on the external surface of the
lower part of the extremities, as in
not getting to the internal parts, provided
by the person, as wounds made by the
state of the health of the person &c

Superficial Ulcers most dangerous than
in common, not all ulcers ought to be
healed, as for instance in gonorrhea
it is often better an abscess burst or
a disorgan has been removed rather than
to let a few months Consumption be
the result.

There are of 3 classes - 1 Hereditary
2 Unhealthy - 3 Specific -

1 Hereditary Ulcers,

30 days of the specific can cure them
The healthy Ulcers throw out white
matter like pain, red pimples &
then the sore can generally be cured
in 10 days. And generally the Ulcers
are like the simple ones but more
or less dry.

then for the simple Ulcers I bring
them together but not tightly ad-
hering that is vomiting.

Of unhealthiness I have seen several
of which I believe belongs, but then
it is not always the same.

[illegible]

Next the Unhealthy Element
of old standing, something like
an indication of Indolent
and in these cases the same
stimulating steam of warm water
from Niagara applied
of course, but a few days
after use in the disease

especially that of a hog, and the
by case of Caribbea. The
leaves are - Cyamum papyriferum
and salt in warm water, A decoction
of stewartia bark, the
the oil of Turpentine mixed with
the essential oil of Camphor, the
and these are very useful
of excruciating complaints, Scirrhus tumour
of deep seated cancer to pain as if pain
when forced in the part as if knocking
the most affecting part of the body
as the neck and breast, and testicles
of males and generally about 40 or
50 years of age - but not easy to dis-
tinguish it but generally has no hard
surface very hard, then bluish cast
the tumour with small portions of fat
particular on the breast and then green
Scirrhus of the breast first begins as a small
lump is felt in the breast and may
taken up in the fingers, but after infla-
mation it adheres to the surrounding
muscles and the gland on the breast
now become affected - the edge is en-
hard on feeling the pain is increased
It may sometimes be got away by

ulcerative process, and where an
Operation is to be performed it to be
done as soon as possible &c
Scrophulous. These may resemble a cancer
but does not show out alike does not
run so. Much, the scirrhous tumour
is small and of a bluish hue.

The fungus hematois is generally of
soft tissue & parts which ~~never~~
cancer never affects,

Of the Arterial cause of the Affec-
tion, Dr Gibbon thinks that most of this
affection are caused by small

Arterial
Cause of Cancer, some say it may be cured
by internal remedies and local applica-
tion - as phosphate of Iron &c but
some say it cannot be cured without the
knife which be considered more effectual
and less pain and then cut it some
distance into the sound flesh

Cancer of the eye not frequent, first appear-
ance in the Cornea, the only remedy for
this is taking out the eye, and some
times it turns -

Cancer of the Lips, there are three sort
it is generally to be about the size
of a shot of a bluish colour, some
times as large as a nut or egg be-
fore it breaks out - sometimes before
it breaks out, the lymphatic glands
are affected and under these
circumstances they never recover
and never when operate on
the lower lip generally affects the upper
lip too, and the kind of small white
appearance - These mutations
are different appearances but little
the but course of scars to be
seen in a different way - sometimes
form of Snow, Canceria the best
milder canidies always increase
it - And cannot be cured in the
advanced stages of the disease
if the lymphatics are not affected the
Cancer may be ~~affected~~ cured by
Operation and even a small pain
course and the tumour cut out
and a little care left him and not
take care the scar too soon and
some Regeneration, the lower section
is more the lips -
The eye is cured as well mutation
the tongue affected by the disease

organs are lined by epithelium
where the tongue is cracked and
swelled so large as to almost pro-
duce suffocation and to be kept
well into the teeth by bandages
about the head &c
it can arise like a tongue
an affection of virus as from
rubbing it against the teeth
or ragged one and this is to be
filled & smooth and is generally
is cured - Operation of the tongue
is best to apply a ligature
around it and attempt it with the
knife &c

Cancer of the breast and the female
more subject to Cancer and the
tumour next - Commence in a
manner - the nipple becomes cracked
and become black and some-
times it is large & hard and more
easily cured than when small -
appears in a generally about the
areola pit or between the nipple &
distinguished from other tumours
as it becomes hard like a lump of
the most certainly the irregularity
of the surface & immense it can
usually if it moves about the

may easily operate upon the
if the secret ^{is} ~~hard~~ ^{is} ~~you~~ ^{is} ~~more~~
not expect so much success for
lumps sometimes happen in are called
milk lumps or sometimes called
in Catamenia which when cured
the lumps go away —
~~Lentamente~~ put a plaster over the
breast and treat the general
system and do this when you do not
suppose it to be a Cancer. Be-
in all operation of the breast the soft
cancer is as soon as you cut them
Cancer of the Uterus next in frequency
to that of the breast — Polypus sometimes
when the finger is introduced into the
vagina. A very hard lump is felt and
when cut with a knife with is covered
with blood and offensive matter it
is very painful — There are 2 varieties
of this disease the one never ulcerates
but the other there is much discharge
and the neck is not much affected
and it must be kept very clean. De-
fecting ~~Polypus~~ ^{Acids} ~~Polypus~~ ^{Polypus}
Inflaming, when the neck only
is affected it may be operated
on by drawing down the uterus

and cut off the neck by a curved
knife & stopping the hemorrhage
by sponging, & then the patient is
healed. —

Cancer of the Rectum seldom
among men and common among
women of the lower, the living mem-
brane of the gut become thick by
the adjacent glands become
affected, the whole system is affec-
ted, some time blocked up and this hap-
pens sometimes before we can see
the disease, when it is curable.
There is no cure, we can only relieve
the symptoms by introducing an
Opium pill into the rectum occa-
sionally and other palliating re-
medies — we must sometimes introduce
a bougie or sponge cover with
Wax &c — Cancer of the penis
which we believe often mistaken for
other disease. The benignant
appearance very much of the soft
Mushroom — But the Cancer is
different having a large hard
deep lacinated base sometimes
high up in the body and cannot

be operated upon with success
in this last stage &c
When the Penis is to be cut off
do not cut it off in one cut
but to hold the penis naturally
cut gradually still holding
the penis and tie up the artery
as you go on which are
haps 6 or 8 no occasion to
introduce any instrument
to carry a hose to Urine
Cancer of the Testicles. When
it is any other disease than is now
of that deep seated lancinating pain
which is common in Cancer. There
are a great number of diseases
which we believe is often mistaken
for Cancer - Different Mode of
Operation which will be shown
another disease is called Fungus or
hematomas, which is a species of
Cancer resembles the substance of
the brain, and more vascular
than Cancer - And often attacks the
liver often the pancreas but
Cancer most attack

Gonorrhea a large leaver poultice
and continued as long as it agrees
especially at night like during this
a milk injection, or sugar of lead
solution of Nitrate of Silver after
it has been continued some time
give a bougie and introduce it
into the Urethra & another injec-
tion of decoct. of Camomile flower
sometimes, Whisky and Water &c
and best Curac Cortic Adulter-
ed Nothing so good for Chordee as
Champhor alone or Combined
5 or 6 grs at bed times &c
hernia puerilis cured by pump-
ing and bleeding and supporting
the parts by bags of Serice & Quina
sometimes, draw away the enlarged
testicles or tape leeching
Women generally do not suffer
so much as men but often as very
severe - not easily distinguished
from Fluor albus, but Fluor
albus comes on with pain in
Loins and very slow, but goes
offe come on in a few days

phlegm. co of P^o 2 stages
1st Phlegm comes on parts of the
body less are very subject to it
and this arises from immediate con-
tact - The phlegm generally occu-
pies the glands penis or in the
nabothian, the first appearance
of Chancres much resembles a small
pimple and when let out matter
is clear, but it a more decided ap-
pearance is where the surgeon can
take it up as a piece of Casti-
lone or General foul matter
about the sore and difficult to
be removed & this is the appear-
ance on the glands penis &c.
but when it affects other parts it
is sometimes different appearance
of in other parts & is often de-
scribed of the even the bluish heat
of the Chancres may not be
seen &c.
The Bubo of the Chancres on the right
side the Bubo is on the same side
& the Bubo goes away in the course
of 2 or 3 days, and unless the system
is sometimes at suppuration &c.

the skin have it in throat —

Treatment After 2 or 3 days
after it breaks out — it has existed for
2 or 3 L. Caustic is true applied, if
it has continued for 2 weeks it is not
worth while to apply caustic &c
If compound mercury have used we often
have secondary symptoms come
on the surface of the skin and the
lesions are affected, but give him
Mercury until the gums are affected
a coppered taste in the mouth, loath
white and gums turned &c
where there is bubo give the Mercury
internally, but when there is
no bubo it may be rubbed on the
thigh or femur is the best
the Calomel a gr morning
and evening or the black pill
If you use mercurial ointment
rub a piece of about size of a
nutmeg ^{of black wart} on a grain of Calomel
and lime water — Mercurate of Soda
to old ulcers, the Labig shelled
in old women must be broken

is a cat of hem lock
Bals, Rhois Antiphlogistic
is not to be rubbed by Mercu-
ry, the best application is
blister and cut a hole in
the middle of the blister to
prevent suppuration, when
the matter must come out
make a number of punctures with
a needle.

Syphilis of second stage sometimes the
skin is affected or the tonsils are
affected difficult from the chronic.
Sometimes one tonsil is destroyed and
a great ulceration of throat and
glottis upwards and destroy the
palate some between the nose and
the mouth, shortly after the throat
is affected there are other affection
as red spots about the size of a
dime or larger and then
have venereal spots, after skin
the joints are affected as the ra-
dium, when as it be, the skin
is painful especially when

swollen in head &c this swelling
retains continues for even 2 years
now or less, as the ~~pain~~ is dis-
charged, the small pimple
about the glands penis are
called venereal warts &c
Conglomerate tumours about the
anus not always - tumours form
about the fingers the ends of the
bone of the fingers become enla-
ged &c - Sometimes the throat
not affected, or not till after the
skin - Mercury generally
given in different forms
are preferred to a lozel or plume
pill - Corrosive sublimates is
sometimes to be preferred - some
cannot be preferred - and when
this is the case, after to fumigation
seat him on a chair with a hole
enclosed with a blanket - the
fumigation passing through the hole
in the chair the Mercury is sure to
act - but does not last so long in
this way and this is the objection
to its process - for this purpose

The Simpliciter Mercury is to be
prepared, when you fail with Cal-
omel or plumb pill you resort
to the above.

For the treatment of the Ulcers of
the throat gargles of Corrosive
Sublimate, but where the bone
is affected we must use Swain-
son's &c. Prescrip. to draw
largely of Sarsaparilla during the
time of taking Swainson's &c.
sometimes ~~the~~ it will purge when
this is the case the dose must be
diminished &c. local of the No.

is blisters sometimes successful or an
internal use of arsenic General
blisters Muricated Linctus Iron or
Caustic.

Fracture &c. Bones may be broken
by Muscular Action and is the most
common cause. Signs of fracture very
variable, and we generally find it An-
gular, but may be Chancous or Clean
or Compound, and Circumference that is torn
ning &c. — Fracture are either
Transverse and are most common and
the fibres separate &c.

Longitudinal, as by gun shot wound
of one of the Condyles you be know
of - as also the Tail as by fracture
tooth - Oblique -

Sometimes it is where it is long is made
complicated as where it is compound with
some other such as rupture of an Artery

Fracture do not heal in old persons as
it does in children - the middle person
do not heal in less than a year as the thigh

Where the thigh is broken we must
use extension and Counter Extension
but must use to give extension &c

I object to place up the parts
where there is too much swelling or not
to receive whether there is a fracture

In this case we are governed by Crep-
itation and is a most certain sign of
fracture of the thigh bone. But the

strips of Stultet's preferable to the
18 tail bandage &c the splints
are of book binders board are prefer-
able and this part in children and

then may be ~~and~~ altered to the limb
Fracture of the foot - There is very profuse
hemorrhage

Fracture of the lower jaw and of the
Upper ~~se~~

vertebrae seldom fracture generally of the
spines process but the skull is broken
remains - fracture of the vertebrae gen-
erally affects the 12th upper spinal small
matron mostly situated the 12th Cervical
and affected, then cut down to the
bone - but draw off the water from the
du de

Of the ribs known by the difficult res-
piration and excitation, difference in
menstruation external or internal in-
jury not much to be done, a collar around
the ribs quite tight

Stomach apt to become carious, and then
perforate and let out the matter, must
sit up, and apply a bandage &c

Of the clavicle simple or Compound signs
very evident head always inclined to the
affected side, bandage &c

If the scapula not often broken, the acro-
mion process is broken and then the arm falls
about as long as the humerus

Fracture of the humerus generally of the
neck not of all other parts if it be
the neck make extension and counter
extension, and then with a roller com-

measure at the finger and hand &
upward. And apply the splints,
External, ^{anterior} internal & posterior and then
a little tow under the splints at the
ends and continue the roller
over the splints, and then a pad as
in fracture of the clavicle secured
by a bandage as the clavicle &
the hand in a sling & —
of the middle of the bone & —

Of the head the bone is often absorbed
& shortened & — of the middle
bandage and apply 4 splints and roller
but no pad, but low as in the neck
the roller about ten or 12 yards long
and 3 inches wide —

Of the Condyle one or both to which
great swelling ensue, the arm to be kept
treatment a bandage so then splints
2 angles or splints and the roller &
tow in the palm of the hand and at
the extremities of the splints

Of fracture of the fore arm and
are very liable to be broken when both
bones are broken they are generally
at the same place

The arrangement generally internal
towards the anterior part, and
mostly takes place at the lower part
of the radius or both often about the
middle - Fracture of the Ulna
appears at the lower part
Treatment for one or both bones the
same if broken - Extension & Con-
traction, two compresses one
about half an inch in thickness to
be placed in the palm of the hand, one
compress internal and external
the ~~two splines~~ a roller next two
splines external and internal and
the roller continued beyond the
elbow - Flexor - summate of
the Ulna Treatment always to be
kept on Extension apply a roller
to be extended higher up - apply tow
at the inside of the elbow and pull
up the halves then apply a spline
over the tow and continue the ro-
ller on

Bones of - Carpus sometimes broken
the turning of a gun &c

When not shattered it is only necessary
to combat inflammation ~~in the~~.

If the fingers are fractured apply
a roller then splint longer than
the fingers and soaked in water &
then continue the roller ~~on~~

~~on~~ And this continue it around
all the fingers and hand.

Fractures of the Pelvis not frequent
happens the Innominata or Sacrum
Sometimes in Old Subject the Os Coccygii
a very little to do not but attend to the
generally Great ment -

Of the Patella often happen by the
too great force of the Extensor Mus-
cle and Fracture are Transverse - obli-
lique or longitudinally.

First apply a short roller the hole limb
of the leg & over this apply a roller
bandage the whole length of the
leg and then a long splint the same
than the leg with two and a band-
age over this the whole length
of the leg &c.

Fracture of the Thigh bone at any
many different places, but
ably about the middle in young
subjects -

In this subject the neck is mostly broken, and sometimes within the Capsular & And here it seldom ever can be cured - but without Capsular ligament it unites as readily as in any other part of the bone, because the ~~it~~ cannot receive sufficient nourishment. If the Proximal is sometimes broken off & in fracture the limb is shortened 2 or 3 inches and is generally standing out - upwards or inwards and will turn on its axis but not so in luxation. In the middle you find a grating

Treatment of Fracture of the Thigh bone first the strips of Scullatins then roll up the long splint in the splint cloth inside and out side the splints of Dr Physick with cushioning between. the splint and leg then apply a handkerchief around the Peroneal and likewise around the foot also around the body ^{and upper part of} with paste board on front and behind. Around all these splints apply a number of pieces of tape &c

An apparatus as improved by Dr
Gibson is where the sound leg acts
as a counter extension to the
one also a bed frame with holes to be
laid on the bed and raised up as
occasion shall require, the bed
should always be hard and not a
feather bed &c

Another something common is more simple
and said to be for experience for
young subjects

Fracture of the Tibia & Fibula
about the ankle

Treated the same as above. The
joint of the leg should be broken ~~the~~ near
the ankle joint the always find
the foot completely turned outward
which is to be treated by bands & d
splints so as to draw it inward
Bones of the foot of Os Calcis
the thigh is bent the leg is bent but
the foot extended then a short
piece from the upper part of the
and continued upwards - the band
is then a bandage serving the
flex position, the same apparatus
for the use of the tibia & fibula

If the bone has fracture of the foot
up a wheel or the shoe may be used
not to bandage up but attend
to Circumference

Compound Fractures & Complica-
ted Fractures where some Artery or
Vein is likewise broken by which
Mortification may take place
In this Case cut down on the
up and that is as a Compound Frac-
ture &c

Amputation is necessary only when
the Arteries Veins Nerves and Mus-
cles are torn and not operated
on immediately especially while
the pulse is low but administer
Anals ~~and~~ danger in Tetan-
us - do not therefore operate
in less than $\frac{1}{2}$ an hour or one hour
In Tetanus the best give plen-
ty of Calomel &c

Dislocations of the thigh bone which
are 4 different upwards and
backwards is the most common
Dislocation of the thigh do not often

rapper, legs are the foot turned
inwards and leg shortened 2 or 3
lines. In Fracture foot is turned
outwards &c

Downwards & forwards next in
frequency; lodged in foramen Ovale
foot turned outwards and leg is
lengthened. Make extension &
Counter Extension secure the Pel-
vis by a band over it

Patella luxated outwards genera-
ly extends the leg on the thigh and
force it in by the fingers of the
surgeon. Person generally
subject it Knocked Kneecap.

Of the Knee joint almost always
followed to death unless ampu-
tated as 'all the ligaments
are torn up

Ankle joint generally with
a fracture

Luxations of the Clavicle which
very seldom happens, but may take place
either at the humeral or sternal extremit-
ties or even both may happen, when the
happens apply the bandage of Jefferson
or for Fracture &c

Luxation of Ribs apply a tight bandage
and low diet &c

Luxations of the Spine

Luxation of the Os Humerus may hap-
pen in 3 different ways ^{upward} ~~upward~~, down-
wards & ~~backward~~ ^{backward}. Signs is a hollow
under the Acromion process and not able
to describe a Cone with the hand &c
Cannot easily happen backward
It may also be of a compound Frac-
ture. Treatment &c

Luxation of the Elbow with the
humerus in 2 ways ^{upward} ~~upward~~, down-
wards & backward

Luxation of Radius &c - posterior
may happen at the anterior part &c
By the loss of rest break the Magna
of the fingers & thumb

Causes of the Spine by distortion in
the cartilage and bag of spine
by which back is bent & injured
which pain in the back and part
of stomach and legs become weak
and paralyzed and with
difficulty support the head very
easily fall —

Remedies are the
forming issues on the back with
administering laxatives daily on
every other day thereby forming
having a continual discharge
exercise carefully taken &
hip joint disease, first
whiptornis pains inside or out
side of the knee joint pain
on feeting the knee Treatment
as above to establish an issue
given, Exercise daily keeping
the limb extended if too weak
decumb up never let out the
matter out allow it to come
spontaneously if it can be pre-
vented, —

* Causes of the Spine. Metastasis or
Carpal bones never amputate as con-
sumption must follow but apply some
Wild Acid, or Arsenic with full dose

Anchylosis a stiffening of joint
sometimes the termination of a for-
midable disease as a Rheuma-
tism or hip joint which is
called a complete Anchylosis but
another known by incomplete
the Complete Anchylosis never to
be touched or attempted to be broken
up but the incomplete ~~they~~ may
be perhaps broken up when
after ^{an} ~~an~~ fracture when we
have not been careful to move
the limb after 4 or 5 weeks to change
the limb from flex to extended
vice versa - When injuries take
place in the Arms we keep it in
the flex position or when it
the leg we keep them in exten-
ded position &c

Treatment we use friction with
Oil or warm water friction to
gradual from then may be the
greatest benefit, but nothing
to be used in consequence of hip
joint &c

Cephalgia or Cartilaginous
formed in joints may form
in the hip Elbow but more com-
monly in the knee generally
arises from straining, often with
this a large quantity of synovial
matter, may sometimes be reliev-
ed by wearing a Cap which will
prevent the Cartilage of the joint
from moving &c.

alluvion, when convenient we re-
move it from the inside of the
condyles, perhaps not best to close
the wounds too soon &c. in
danger of Inflammation or Tetanus

Distortion or Club foot, or wound
shoulder or elbow or spine or head
may be twisted by some awkward
position, most common Club
foot

Gonarthrosis or Disparity of the knee
joint more common to old farri-
ng men, follows Typhus fever or Rheuma-
tism, by blistering or leeches let out
the blood, make use of a bandage

White Swelling generally
arises from cold between 30 &
40 years of age

Rheumatic white swelling or
Inflammation of the Synovial
generally affects the Knee joint
Morbic enlargement of the Syno-
vial Membrane subjected to
Children pain in both comes
on gradually.

Scrophulous White ~~Children~~ Swelling
in Scrophulous Children upper
extremity of the Tibia enlarged
Exostosis happens to long bones
Circumscribed, Spicuous, lami-
lated & Tuberculated exostosis

With lamellated exostosis is joined
and necrosis

Exostosis is bone matter thrown
out, not after necessity to re-
move, but it sometimes becomes
necessary


Osteo sarcoma perhaps a can-
crous affection of the bone &
very difficult to cure


Osteo ventosa An Expansion of
the bone a shell of the periosteum
Necrosis a decay of the bone


Dec
Diseases of the Arteries


Aneurysm not a dilatation
of the Arteries but a rupture of
the internal and middle &c


Treatment of White Phlegm
patient to be placed on a hard
mattress and apply a ~~str~~ straight
joint carved in in shape to pre-
vent motion - then purge 2 or
3 times a week or after if
the patient will allow of it and
then make issue ~~over the~~















MSS 2/0213-01

Acc. 1991-086-01